


B34



(11) Publication number:

**0 400 661
A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90110399.4

(51) Int. Cl.⁵: **C07D 403/04, C07D 409/04,
C07D 409/14, C07D 239/42,
A61K 31/505**

(22) Date of filing: 31.05.90

(30) Priority: 01.06.89 US 360657
30.03.90 US 503197

(43) Date of publication of application:
05.12.90 Bulletin 90/49

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: **BRISTOL-MYERS SQUIBB
COMPANY**
345 Park Avenue
New York, N.Y. 10154(US)

(72) Inventor: Yevich, Joseph P.
115 Crest Road
Southington, Connecticut 06489(US)
Inventor: Dextraze, Pierre
6600 Place de la Bataille
La Prairie, Quebec(CA)

(74) Representative: Kinzebach, Werner, Dr. et al
Patentanwälte Reitstötter, Kinzebach und
Partner Sternwartstrasse 4 Postfach 86 06 49
D-8000 München 86(DE)

(54) Agents for treatment of brain ischemia.

(57) A series of 5-halopyrimidin-2-ylpiperazinylalkyl derivatives having useful anti-ischemic properties for treatment and prevention of disorders resulting from brain and/or spinal cord anoxia.

EP 0 400 661 A1


BEST AVAILABLE COPY

AGENTS FOR TREATMENT OF BRAIN ISCHEMIA

BACKGROUND OF THE INVENTION


This invention pertains to heterocyclic carbon compounds having drug and bio-affecting properties and to their preparation and use. In particular, the invention is concerned with 1,4-disubstituted piperazine derivatives wherein one substituent is a 5-halopyrimidin-2-yl moiety and the other is a carbon chain bearing a carbocycle or heterocycle moiety at its terminus, usually via a linking hetero atom or functional group moiety.

Related art may be viewed in light of the following general structural Formula 1.




in which Ar is an aromatic ring, usually phenyl; X is a carbonyl or carbinol group; alk is an alkylene chain.

20 The most closely related art would appear to be U.S. 4,605,655, issued to Yevich, *et al.*, on August 12, 1986. This patent disclosed and claimed piperazinylbutyrophenone derivatives possessing neuroleptic properties and characterized by structural Formula 2.




wherein X is




With R being C₁₋₄ alkyl, hydrogen or fluorophenyl; and Y is hydrogen or halogen. The instant anti-ischemic compounds are distinguished from these art compounds either by the nature of the terminal carbocyclic ring system, the nature of X, the presence of an alkylene bridge on the piperazine ring and the nature of the R³ substituent.

40 In U.S. 2,973,360, issued February 28, 1961, a series of CNS depressant compounds is disclosed with Ar being 2-thienyl; X being carbonyl or carbinol; and alk being C₂ and C₃ alkylene. The most pertinent compound specifically exemplified and claimed in this patent is shown below as structure (3).




50 The following references, while related, are felt to be less relevant to the new compounds disclosed in this application.

Reginer, *et al.*, U.S. Pat. No. 3,299,067, issued January 17, 1967 discloses compounds comprising a benzyl-type moiety attached to the 2-pyrimidinylpiperazine. A specific example of this series which is said to be useful as peripheral vasodilators, analgesics and antiinflammatory agents, is shown below as structure (4).




(4)

U.S. Pat. No. 3,802,210 issued to Regnier, et al., in April 1974 relates to a series of aryloxypropanolamine antihypertensive compounds having a pyrimidinylpiperazine moiety as in (5). However, these compounds are not butyrophenones or close analogs.




(5)

U.S. Pat. No. 4,316,899 issued to Markwell on February 23, 1982 relates to another series of aryloxypropanolamine antihypertensive compounds containing a pyrimidinylpiperazine moiety as exemplified by structure (6)




(6)

Gotti, et al., JPET, 247'3, pages 1211-1221 (1988); have disclosed that ifenprodil and a derivative are effective in tissue sparing in animal models of stroke and brain infarction.




Wauquier, et al., in "Drug Development Research", 8:373-380 (1986) disclosed that Sabeluzole (R 58,735) is a potent antihypoxic agent with anticonvulsant properties.



A series of anti-anoxic 2-[4-benzoyl-1-piperidinyl]-1-phenylalkanol derivatives, having some structural resemblance to ifenprodil type compounds, is disclosed in U.S. 4,711,899 issued in December, 1987 to Gaudilliere, et al.


There is nothing in these references, or in the general prior art, to suggest the anti-ischemic compounds of the present invention.

In its broadest aspect, the present invention is concerned with 5-halopyrimidin-2-yl piperazine derivatives having anti-ischemic properties characterized by a compound of Formula I




wherein

Z is a member selected from the group consisting of



20 naphthalenyl, anthracenyl, fluorenyl, phenanthrenyl, and C₅₋₆ cycloalkyl. X is a member selected from the group consisting of -O-, -S-, -SO₂-, -CO-,

-CH(OH)-, wherein R⁴ is hydrogen, C₁₋₄ alkyl, and -CH(R⁵)- wherein R⁵ is hydrogen, CN, or NHR⁶ with R⁶ being acetyl,



30 wherein W is hydrogen, halogen or alkoxy;
or Z and X taken together can be



R¹ is hydrogen or C₁₋₄ alkyl; R² is halogen; and R³ is hydrogen, C₁₋₄ alkoxy or C₁₋₄ alkylthio. The symbol n is the integer 1 - 3 and the symbol m is the integer 0 or 1. There is a proviso for the compounds of Formula I which is that Z cannot be when X is

40 -CH(OH)- or -CO- while R³ is either hydrogen or C₁₋₄ alkoxy, or while m is 0.

It is to be understood that pharmaceutically acceptable salts and/or solvates of the Formula I compounds also comprise the present invention. Further, as used herein, halogen denotes chlorine, bromine, iodine and preferably fluorine. Preferred compounds are those wherein Z is p-fluorophenyl and wherein X is -CH(R⁵)-.

45 It is also to be understood that the present invention is considered to include stereoisomers as well as optical isomers, e.g. mixtures of enantiomers as well as individual enantiomers and diastereomers, which arise as a consequence of structural asymmetry in selected compounds of the present series. Separation of the individual isomers is accomplished by application of various methods which are well known to practitioners in the art.

50 For medicinal use, the pharmaceutically acceptable acid addition salts, those salts in which the anion does not contribute significantly to toxicity or pharmacological activity of the organic cation, are preferred. The acid addition salts are obtained either by reaction of an organic base of structure I with an organic or inorganic acid, preferably by contact in solution, or by any of the standard methods detailed in the literature available to any practitioner skilled in the art. Examples of useful organic acids are carboxylic acids such as
55 maleic acid, acetic acid, tartaric acid, propionic acid, fumaric acid, isethionic acid, succinic acid, pantoic acid, cyclamic acid, pivalic acid and the like; useful inorganic acids are hydrohalide acids such as HCl, HBr, HI; sulfuric acid; phosphoric acid and the like. The preferred solvate forms of Formula I compounds are

hydrates.

The compounds of the present invention are useful pharmacologic agents with anti-ischemic properties. Brain cells are particularly vulnerable to damage caused by ischemic conditions. Brain ischemia, or insufficient oxygen, may result from injury or disease and may last from only transient periods of time to periods of lengthy duration, as in stroke. In this regard, the compounds of Formula I are useful for treatment and prevention of injury to the brain and spinal cord and of edema due to head trauma, stroke, arrested breathing, cardiac arrest, Reye's syndrome, cerebral thrombosis, embolism, hemorrhage or tumors, encephalomyelitis, spinal cord injury, hydroencephalus, and post-operative brain injury.

The anti-ischemic activity of the compounds of Formula I have been demonstrated in certain pharmacologic tests and model systems that are used to determine drug effects on brain ischemia and its aftermath. Most specifically, administration of the compounds of Formula I results in protecting against hypoxia-induced death in anoxic nitrogen test in rats. This particular test identifies the neuro-protective effects of substances against lethal brain damages produced by a lack of oxygen consumption (anoxia). In this test procedure, control animals exposed for one minute to a pure nitrogen atmosphere will expire because of respiratory failure caused by irreversible damage to the brain respiratory center. The animals exhibit strong heartbeat following anoxia exposure. To demonstrate effectiveness, experimental compounds must antagonize the anoxic insult resulting in survivability of the test animals.

One aspect then of the present invention involves administration of a compound of Formula I or a pharmaceutically acceptable acid and/or solvate thereof, to a mammal suffering from ischemia or being susceptible to ischemia. In general the compound would be given in a dose range of from about 0.1 mg/kg to about 10 mg/kg body weight.

Although the dosage and dosage regimen of a Formula I compound must in each case be carefully adjusted, utilizing sound professional judgment and considering the age, weight and condition of the recipient, the route of administration and the nature and extent of the ischemia, generally, the daily dose for human use will be from about 0.1 g to about 10 g, preferably 0.5 g to 5 g, when given orally. In some instances, a sufficient therapeutic effect can be obtained at lower doses while in others, larger doses will be required. As is apparent to one skilled in clinical pharmacology, the amount of a Formula I compound comprising the daily dose may be given in a single or divided dose, taking into account those principles understood by the skilled practitioner and necessary for his practice of the art.

The term "systemic administration" as used herein refers to oral, sublingual, buccal, transnasal, transdermal, rectal, intramuscular, intravenous, intraventricular, intrathecal, and subcutaneous routes. Generally, it will be found that when a compound of the present invention is administered orally a slightly larger quantity of the active drug may be required to produce the same effect as a somewhat smaller quantity when given parenterally. In accordance with good clinical practice, it is preferred to administer the instant compounds at a concentration level which will produce effective beneficial effects without causing any harmful or untoward side effects.

Therapeutically, the instant compounds are generally given as pharmaceutical compositions comprised of an effective ischemia-protective amount of a Formula I compound or a pharmaceutically acceptable acid addition salt and/or hydrate thereof and a pharmaceutically acceptable carrier. Pharmaceutical compositions for effecting such treatment will contain a major or minor amount (e.g. from 95% to 0.5%) of at least one compound of the present invention in combination with a pharmaceutical carrier, the carrier comprising one or more solid, semi-solid, or liquid diluent, filler and formulation adjuvant which is non-toxic, inert and pharmaceutically acceptable. Such pharmaceutical compositions are preferably in dosage unit forms; i.e., physically discrete units having a pre-determined amount of the drug corresponding to a fraction or multiple of the dose which is calculated to produce the desired therapeutic response. In usual practice, the dosage units contain 1, 1/2, 1/3, or less of a single dose. A single dose preferably contains an amount sufficient to produce the desired therapeutic effect upon administration at one application of one or more dosage units according to the pre-determined dosage regimen, usually a whole, half, third, or less of the daily dosage administered once, twice, three, or more times a day. It is envisioned that other therapeutic agents can also be present in such a composition. Pharmaceutical compositions which provide from 0.1 to 1 g of the active ingredient per unit dose are preferred and are conventionally prepared as tablets, lozenges, capsules, powders, aqueous or oily suspensions, syrups, elixirs, and aqueous solutions. Preferred oral compositions are in the form of tablets, capsules, and may contain conventional excipients such as binding agents (e.g., syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone), fillers (e.g. lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine), lubricants (e.g. magnesium stearate, talc, polyethylene glycol or silica), disintegrants (e.g. starch) and wetting agents (e.g. sodium lauryl sulfate). Solutions or suspensions of a Formula I compound with conventional pharmaceutical vehicles are employed for parenteral compositions such as an aqueous solution for intravenous injection or an oily suspension for intramuscular injection. Such


compositions having the desired clarity, stability and adaptability for parenteral use are obtained by dissolving from about 0.1% to 10% by weight of a Formula I compound or one of its salt forms in water or a vehicle consisting of a polyhydric aliphatic alcohol such as glycerine, propylene glycol, and the polyethylene glycols or mixtures thereof. The polyethylene glycols consist of a mixture of non-volatile, usually liquid, polyethylene glycols which are soluble in both water and organic liquids and which have molecular weights from about 200 to 1500.

When transnasal application is intended, the Formula I compound pharmaceutical composition is formulated in a pharmaceutical composition which enhances penetration of the nasal mucosa. Such formulations normally employ fatty acid salts of the Formula I base compound and their preparation and use would be known to one skilled in the pharmaceutical arts.

The general procedures for preparation of Formula I compounds are outlined in Schemes 1 and 2. In these schemes the symbols A and B refer to subclasses of the moieties denoted by X supra.

Scheme 1

General Synthesis of Leading to Formula I Compounds



In scheme I: Z, R¹-R³, m and n are as previously defined. The symbol A represents O, S, CO and CH₂ and symbol W represents a leaving group, as well understood in organic synthesis, such a leaving group being preferably chloride or bromide. In addition Z and A can be taken together as a ketal such as




The intermediate compounds II and III are known in the literature but some synthetic examples will also be given hereinbelow for convenience in preparing the product compounds of the present invention.

In particular, synthesis of Formula II intermediates as well as some IA analogs are disclosed in U.S. 4,605,655 which is incorporated herein in its entirety. As covered in U.S. 4,605,655, when preparing IA wherein A is CO, the carbonyl group is generally protected as the ketal derivative.

Scheme 2 outlines chemical conversions of IA product compounds to IB product compounds by transformation of certain X moieties (IA-->IB).


Scheme 2

Variation of X in Formula I Compounds



Transformations

<u>A</u>	<u>Reaction</u>	<u>B</u>
S	oxidation, e.g. H_2O_2	SO_2
CO	reduction, e.g. $NaBH_4/EtOH$	CHOH
CO	reductive alkylation e.g. R^4MgCl	CR^4OH
CO	Tosylmethyisocyanide	CHCN
CHOH	$Ph_3P/(i-PrO_2C-N)_2/(PhO)_2PON_3$	N_3
N_3	cat. H_2	NH_2
NH_2	acylation or sulfonylation	NHR^6
<u>Z-A</u>		<u>Z-B</u>



In scheme 2, Z, R^1 - R^3 , A, n and m are as previously defined and symbol B represents other values of X such as SO_2 , CR^4OH and NHR^6 . Additionally B can be N_3 and NH_2 , thereby giving 2 synthetic intermediates in the process of converting $X = CHOH$ to $X = CHNHR^6$ in Formula I compounds.

DESCRIPTION OF SPECIFIC EMBODIMENTS

The compounds which constitute this invention and their methods of preparation will appear more full from a consideration of the following examples which are given for the purpose of illustration only and are not to be construed as limiting the invention in sphere or scope. All temperatures are understood to be in degrees C. when not specified.

The nuclear magnetic resonance (NMR) spectral characteristics refer to chemical shifts (δ) expressed as parts per million (ppm) versus tetramethylsilane (TMS) as reference standard. The relative area reported for the various shifts in the proton NMR spectral data corresponds to the number of hydrogen atoms of a particular functional type in the molecule. The nature of the shifts as to multiplicity is reported as broad singlet (bs), singlet (s), multiplet (m), doublet (d), doublet of doublets (dd), or quartet (q). Abbreviations employed are DMSO- d_6 (deuterodimethylsulfoxide), $CDCl_3$ (deuteriochloroform), and are otherwise conventional. The infrared (IR) spectral descriptions include only absorption wave numbers (cm^{-1}) having functional group identification value. The IR determinations were employed using potassium bromide (KBr) as diluent. The elemental analyses are reported as percent by weight.

SYNTHESIS OF INTERMEDIATES

EXAMPLE 1

γ -Chloro-p-fluorobutyrophenone ethylene ketal

A solution of γ -chloro-p-fluorobutyrophenone (50 g, 0.25 mole, commercially available); ethylene glycol (50 mL); p-toluene sulfonic acid (0.1 g) in 300 mL benzene is refluxed for 18 h with water of reaction being removed by means of a Dean-Stark water trap. Upon cooling to room temperature, the reaction mixture is washed with dilute sodium bicarbonate, dried (MgSO_4), filtered and the benzene removed by concentration in vacuo. The residual oil was distilled to give 57.7 g (93%) of product, b.p. 106-112.0.01 Torr.

EXAMPLE 2

1-(5-Fluoro-2-pyrimidinyl)piperazine

(1) Ethyl 4-(5-fluoro-4-methylthio-2-pyrimidinyl)-1-piperazine carboxylate: A mixture of 2-chloro 5-fluoro-4-methylthiopyrimidine (28.3g, 0.16 mole), N-carbethoxypiperazine (25.26 g, 0.16 mole), anhydrous K_2CO_3 (66.0 g) and a catalytic amount of KI in MeCN (400 mL) was stirred and heated under reflux for 18 h. The hot reaction mixture was filtered, concentrated in vacuo and the residue crystallized from EtOH to give 29.8 g (62%) of intermediate product.

(2) Ethyl 4-(5-fluoro-2-pyrimidinyl)-1-piperazine carboxylate: A mixture of ethyl-4-(5-fluoro-4-methylthio-2-pyrimidinyl)-1-piperazine carboxylate (29.8 g 0.1 mole) and Raney Nickel catalyst (15 tsp) in EtOH (550 mL) was stirred and heated under reflux for 48 h. The reaction mixture was filtered, concentrated in vacuo and the residue recrystallized twice from EtOH to provide 11.2 g (45%) of product, m.p. 104-107 °C.

A solution of this ester intermediate (11.2 g, 0.04 mole) in 6N HCl (100 ml) was stirred and heated under reflux overnight. The cooled reaction mixture was made alkaline by addition of 50% NaOH, extracted with Et_2O and the extract dried (MgSO_4) and concentrated in vacuo to provide 7.23 g (100%) of product as a viscous oil which was treated with ethanolic HCl in EtOH to yield the hydrochloride salt, m.p. 250-252 °C.

Anal. Calcd. for $\text{C}_8\text{H}_{11}\text{FN}_4\cdot\text{HCl}$:	
Found:	C, 43.95; H, 5.54; N, 25.63. C, 44.23; H, 5.57; N, 25.38.

EXAMPLE 3

5-Bromo-2-(1-piperazinyl)pyrimidine

To an ice-cooled solution of 1-(2-pyrimidinyl) piperazine (16.4 g, 0.1 mole) in 1N HCl (100 mL) was added dropwise bromine (15.98 g, 0.1 mole). After stirring at 0° for 0.5 h, the mixture was heated to 100 °C until dissipation of the red color had occurred. The mixture was filtered, cooled, made alkaline with 50% NaOH and extracted with Et_2O . The dried extract (MgSO_4) was concentrated in vacuo to provide 14.5 g (62%) of product, m.p. 73-75 °C.

By appropriate modification of this procedure the 5-chloro intermediate and the 5-iodo intermediate may be prepared.

EXAMPLE 4

1-(4-Fluorophenyl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl]butanone hydrochloride (IA)

A mixture of 1-(5-fluoro-2-pyrimidinyl)piperazine (7.3 g, 0.04 mole), γ -chloro-p-fluorobutyrophenone ethylene ketal (14.5 g, 0.06 mole), anhydrous K_2CO_3 (24.8 g) and a catalytic amount of KI in MeCN (100 mL) was stirred and heated under reflux for 36 h. The hot mixture was filtered, concentrated in vacuo and the residue treated with 20 mL of 3N HCl and 100 mL EtOH. After cooling in ice, the product was collected by filtration and dried to give 7.6 g (50%) of product as a white solid, m.p. 234-236 °C.

Anal. Calcd. for $C_{18}H_{20}F_2N_4O \cdot HCl$:	
Found:	C, 56.48; H, 5.53; N, 14.64.
	C, 56.27; H, 5.52; N, 14.27.

1H NMR (DMSO- d_6): 2.10 (2,m); 3.20 (6,m); 3.54 (4,m); 4.58 (2,m); 7.34 (2,m); 8.08 (2,m); 8.55 (2,s); 11.60 (1,bs).

IR (KBr): 960, 1235, 1245, 1365, 1510, 1560, 1600, 1680, 2550, and 2920 cm^{-1} .

EXAMPLE 5

1-(4-Fluorophenyl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl]-butanol hydrochloride (IB)

A mixture of the Ia compound prepared above in Example 4 (7.6 g 0.02 mole) and $NaBH_4$ (2.3 g, 0.06 mole) in EtOH (650 mL) was stirred overnight. The mixture was treated with ethanolic HCl, stirred at room temperature for 1.5 h. then heated to reflux. Solvent was removed in vacuo and to the residue was added 1N NaOH and CH_2Cl_2 . The organic layer was separated, dried ($MgSO_4$) and concentrated in vacuo. This residue was dissolved in EtOH (treated with ethanolic HCl and cooled). The hydrochloride salt was collected by filtration and dried to afford 6.2 g (81%) of product, m.p. 236-238 °C.

Anal. Calcd. for $C_{18}H_{22}F_2N_4O \cdot HCl$:	
Found:	C, 56.18; H, 6.03; N, 14.56
	C, 55.98; H, 6.06; N, 14.23

1H NMR (DMSO- d_6): 1.71 (2,m); 3.10 (4,m); 3.47 (4,m); 4.59 (3,m); 5.30 (1,bs); 7.11 (2,m); 7.40 (2,m); 8.53 (2,s); 11.50 (1,bs).

IR (KBr), 955, 1220, 1235, 1370, 1440, 1455, 1480, 1510, 1560, 1605, 2600 and 2920 cm^{-1} .

EXAMPLE 6

1-[3-[2-(4-fluorophenyl)-1,3-dioxolan-2-yl]propyl]-4-(5-fluoro-2-pyrimidinyl)piperazine hydrochloride (IA)

(1) A mixture of the γ -chloro ketal (Ex. 1: 27.49 g, 0.112 mole), piperazine (48.24 g, 0.56 mole), K_2CO_3 (46.43 g, 0.33 mole), and a catalytic amount of KI in 358 mL of MeCN was refluxed for 18 h. The hot reaction mixture was filtered and the filtrate concentrated in vacuo to a residue which was partitioned between water (250 mL) and Et_2O . The water layer was extracted further with Et_2O , the extracts combined and dried ($MgSO_4$) and concentrated in vacuo to give 28.5 g of 1-[3-[2-(4-fluorophenyl)-1,3-dioxolan-2-yl]-

propyl] piperazine.

(2) This piperazine intermediate (7.8 g, 0.026 mole), 2-chloro-5-fluoro-4-methylthiopyrimidine (4.73 g, 0.026 mole), pulverized K_2CO_3 (11.05 g) and a catalytic amount of KI in 80 mL MeCN was refluxed 18 h. The hot reaction mixture was filtered and the filtrate was concentrated in vacuo to give 11.1 g of residue which was flash-chromatographed on silica gel (3% MeOH:CH₂Cl₂). Appropriate fractions were combined, dissolved in 10 mL EtOH, chilled and treated with ethanolic HCl from which 1.5 g of 1-[3-[2-(4-fluorophenyl)-1,3-dioxolan-2-yl]propyl]-4-(5-fluoro-2-methylthio-2-pyrimidinyl)piperazine hydrochloride, m.p. 233-235° C was obtained.

Anal. Calcd. for C ₂₁ H ₂₆ F ₂ N ₄ O ₂ S.HCl:	
	C, 53.33; H, 5.75; N, 11.85.
Found:	C, 53.53; H, 5.81; N, 12.03.

(3) 1-[3-[2-(4-fluorophenyl)-1,3-dioxolan-2-yl]propyl]-4-(5-fluoro-2-methylthio-2-pyrimidinyl)piperazine hydrochloride (7.45 g, 0.017 mole), triethylamine (3.05 g, 0.034 mole) and 2 teaspoons of Raney Nickel in water were mixed in EtOH (125 mL) and refluxed 18 h. The hot reaction mixture was filtered and the filtrate was concentrated in vacuo to about 15 volume. A crude crystalline product was obtained by filtration and its recrystallization from 20-25 mL EtOH gave 1.6 g of solid, m.p. 220-222° C. This solid was converted to the hydrochloride salt in EtOH using ethanolic HCl. Filtration and drying gave 1.6 g of product, m.p. 242-244° C.

Anal. Calcd. for C ₂₅ H ₂₄ F ₂ N ₄ O ₂ .HCl:	
	C, 56.27; H, 5.90; N, 13.12.
Found:	C, 56.12; H, 6.06; N, 12.90.

EXAMPLE 7

Preparation of 2-(5-fluoro-2-pyrimidinyl)-(1S,4S)-2,5-diazabicyclo[2.2.1]heptane

(1) Trans-4-hydroxy-1-(4-toluenesulfonyl)-L-proline. To a solution of hydroxy-L-proline (80 g, 0.61 mole) in 2N NaOH (800 mL) was added tosylchloride (136.1g, 0.71 mole) in Et₂O (700 mL). The reaction mixture was stirred at 0° C for 1.5 h and continued for an additional 3.5 h at 23° C. The aqueous layer was separated, acidified with concentrated HCl to pH 1 and allowed to stand at -10° C for 12 h. The precipitate was filtered, washed with cold water, and concentrated in vacuo to a volume of 300 mL. The precipitate obtained was combined with the previous precipitate, and the combined solids were recrystallized from ethylacetate. Drying in vacuo at 50° C for 24 h afforded trans-4-hydroxy-1-(4-toluenesulfonyl)-L-proline (107.38 g, 62%).

(2) Potassium salt of trans-4-hydroxy-1-(4-toluenesulfonyl)-L-proline. To a solution of trans-4-hydroxy-1-(4-toluenesulfonyl)-L-proline (107.38 g, 0.376 mole) in acetone (450 mL) was added potassium 2-ethylhexanoate in BuOH (1.91 N; 189.5 mL). After standing at 23° C for 20 min, the insoluble material was filtered and the resulting solution was concentrated to 320 mL. Et₂O (1000 mL) was added to the concentrate and the solvents removed under reduced pressure yielding a solid (122.90 g). The hygroscopic product was used in the next step without further purification.

(3) 1N-Tosylhydroxy-L-proline methyl ester. To a solution of potassium trans-4-hydroxy-1-(4-toluenesulfonyl)-L-proline (122.90 g, 0.376 mole) in 250 mL of N,N-dimethylacetamide was added CH₃I (24.5 mL, 0.39 mole) while under N₂ atmosphere. The light protected mixture was stirred 16 h. The mixture was poured onto ice water and extracted with CH₂Cl₂ (3 x 400 mL). The combined organic extracts were washed with 2% NaHCO₃ (400 mL), H₂O (4 x 1.5 L), dried over MgSO₄, filtered, and concentrated under reduced pressure to leave a viscous oil. The crude oil was triturated with petroleum ether to give N-

tosylhydroxy-L-proline methyl ester as a pale yellow solid (63.20 g, 56.2%) which was used in the next step without further purification.

(4) (2S,4R)-1-(4-toluenesulfonyl)-2-hydroxymethyl-4-hydroxy pyrrolidine. To a solution of N-tosylhydroxy-L-proline methyl ester (62.20 g, 0.21 mole) in THF (600 mL) at 0 °C was added LiBH₄ (15.8 g, 0.73 mole) in small portions. The reaction mixture was stirred at 0 °C for 1 h and allowed to stand at 23 °C for 18 h. The reaction mixture was cooled to -20 °C, made neutral with 6N HCl and concentrated under reduced pressure. The residue was treated with water (550 mL) and extracted with EtOAc (4 x 300 mL). The combined organic extracts were washed with H₂O, dried over MgSO₄, filtered and concentrated under reduced pressure to give N-tosylhydroxy-L-prolinol as a white solid (50.56 g, 88.8%) which was used in the next step without further purification.

(5) (2S,4R)-1-(4-toluenesulfonyl)-2-(4-toluenesulfonyloxymethyl)-4-(4-toluenesulfonyloxy)-pyrrolidine. To a solution of p-toluenesulfonyl chloride (155 g, 0.81 mole) in pyridine (330 mL) at 0 °C was added N-tosylhydroxy-L-prolinol (104.40 g, 0.39 mole). The reaction mixture kept at 6 °C for 72 h and then poured into cold 2N HCl (2.5L). The aqueous layer was extracted with CH₂Cl₂ (3 x 1000 mL). The combined organic extracts were dried over MgSO₄, filtered, and concentrated under reduced pressure to give an oil. The oil was triturated with EtOH and the solid that formed was collected by filtration. The crude product was recrystallized from EtOH (3.5 L) to give tritosylhydroxy-L-prolinol (99.87 g, 44.2%, m.p. 130-132 °C, [α]_D²⁴ = 57.1, c = 1.2, acetone).

(6) (1S,4S)-2-(4-toluenesulfonyl)-5-phenylmethyl-2,5-diazabicyclo[2.2.1]heptane. To a suspension of tritosylhydroxy-L-prolinol (98.87 g, 0.17 mole) in toluene (350 mL) was added benzylamine (54.83 g, 0.51 mole). The resulting mixture was heated at reflux for 18 h and allowed to cool to 23 °C. The reaction mixture was filtered and the solvent removed under reduced pressure. The residue was triturated with EtOH and the solid that formed was collected by filtration to give (1S,4S)-2-(4-toluenesulfonyl)-5-phenylmethyl-2,5-diazabicyclo[2.2.1]heptane (54.18 g, 93.2%) which was used in the next step without further purification.

(7) (1S,4S)-N-benzyl-2,5-diazabicyclo[2.2.1]heptane dihydrobromide. A mixture of (1S,4S)-2-(4-toluenesulfonyl)-5-phenylmethyl-2,5-diazabicyclo[2.2.1]heptane (54.0 g, 0.16 mole) in AcOH (830 mL) containing HBr (30% wt.) was heated at 70 °C for 18 h. The reaction mixture was allowed to cool and concentrated under reduced pressure to a final volume of ca. 300 mL. The precipitate that formed was filtered and washed with acetone to give (1S,4S)-N-benzyl-2,5-diazabicyclo[2.2.1]heptane (50.30 g, 91.3%, m.p. 272-275 °C).

Anal. Calcd. for C ₁₂ H ₁₆ N ₂ ·2HBr:	
Found:	C, 41.17; H, 5.19; N, 8.01. C, 40.83; H, 5.16; N, 8.06.

(8) 2-(tert-butyloxycarbonyl)-5-phenylmethyl-(1S,4S)-2,5-diazabicyclo[2.2.1]heptane. The title compound was prepared as described for the 1R,4R isomer Ex. 8 (10) yield: 8.30 g.

(9) 2-(t-butyloxycarbonyl)-(1S,4S)-2,5-diazabicyclo[2.2.1]heptane. Into a solution of the (8) intermediate (8.30 g, 28.82 mmol) in 250 mL of EtOH was added AcOH (3.2 mL). The reaction mixture was treated with 10% palladium-on-carbon (2.40 g) and hydrogenated at 50 psi for 6 h at 23 °C. The same workup procedure as described for the 1R,4R isomer was followed. Yield 5.60 g, 98.1%. The product was used in the next step without purification.

(10) 2-(t-butyloxycarbonyl)-5-(5-fluoro-4-methylthio-2-pyrimidinyl)-(1S,4S)-2,5-diazabicyclo[2.2.1]heptane (4.50 g, 22.73 mmol), 2-chloro-5-fluoro-4-methylthiopyrimidine [4.64 g (87.5% pure), 22.73 mmol], micropulverized K₂CO₃ (9.40 g, 68.19 mmol), and KI (0.57 g, 3.41 mmol) in 65 mL of MeCN was refluxed for 44 h. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in a minimum amount of H₂O. The solution was extracted with CH₂Cl₂, washed with saturated NaCl solution, dried over K₂CO₃, filtered and concentrated under reduced pressure. Flash chromatography (Hexane EtOAc; 4:1) gave the title compound (6.60 g, 85.4%).

(11) 2-(5-fluoro-2-pyrimidinyl)-(1S,4S)-2,5-diazabicyclo[2.2.1]heptane. A mixture of intermediate (10) compound (6.50 g, 19.12 mmol) and Raney Ni (5 scoops) in 100 mL EtOH was refluxed for 48 h. Raney Ni was filtered through a celite pad and the filtrate was concentrated to give 2-(t-butyloxycarbonyl)-5-(5-fluoro-2-pyrimidinyl)-(1S,4S)-2,5-diazabicyclo[2.2.1]heptane (5.31 g, 94.5%). The product (4.94 g, 16.80 mmol) was dissolved in 3N HCl (100 mL) and refluxed for 3h. The reaction mixture was concentrated under reduced pressure. The residue was made basic with 5N NaOH solution and extracted with CH₂Cl₂ (4 x 100 mL). The combined CH₂Cl₂ extracts were dried over K₂CO₃, filtered, and concentrated under reduced

pressure to give the title compound (2.85 g, 87.4%).

EXAMPLE 8

5

Preparation of (1R,4R)-2,5-diazabicyclo[2.2.1]heptane dihydrobromide

10 (1) Allo-4-hydroxy-D-proline Hydrochloride. A solution of acetic anhydride (380 mL) in glacial AcOH (1.2 L) was heated to 50 °C and 4-hydroxy-L-proline (100 g, 0.76 mole) was added in one portion. The reaction mixture was heated at reflux for 5.5 h. The reaction was cooled and the solvent was removed in vacuo leaving a thick oil. The oil was dissolved in 2N HCl (1.5 L) and heated at reflux for 3 h. The reaction mixture was allowed to stand at 23 °C for 18 h. The solution was heated on a steam bath (~90 °C), treated
15 with activated charcoal, and filtered through celite. The filtrate was concentrated under reduced pressure and the solid was collected by suction filtration to give allo-4-hydroxy-D-proline hydrochloride (107.24 g, 84.2%).

(2) Allo-4-hydroxy-D-proline ethyl ester hydrochloride. A suspension of allo-4-hydroxy-D-proline hydrochloride (106.24 g, 0.63 mole) in absolute EtOH (550 mL) at 0 °C was treated with dry HCl until the reaction became homogeneous. The solution was heated at reflux for 5 h. The reaction mixture was cooled to 23 °C and allowed to stand 16 h. The reaction was cooled to 0 °C the resulting precipitate was collected by suction filtration and washed with acetone to give allo-4-hydroxy-D-proline ethyl ester hydrochloride (95.53 g, 77.5%, m.p. 145-147 °C).

(3) Allo-1-(4-toluenesulfonyl)-4-(toluenesulfonyloxy) -D-proline ethyl ester. To a solution of allo-4-hydroxy-D-proline ethyl ester hydrochloride (101.50 g, 0.52 mole) and triethylamine (52.51 g, 0.52 mole) in pyridine (1020 mL) at -5 °C was added in portions p-toluenesulfonyl chloride (218.10 g, 1.14 mole). The solution was stirred at 0 °C for 1 h and stored at -10 °C for 17 h. The mixture was stirred at 23 °C for 5 h, poured onto ice H₂O (750 mL), and the precipitate filtered. The precipitate was washed with H₂O to give allo-1-(4-toluenesulfonyl)-4-(toluenesulfonyloxy)-D-proline ethyl ester (155.14 g, 62.7%).

30 (4) 4-(Acetyloxy)-1-(4-toluenesulfoxy)-D-proline ethyl ester. A solution of allo-1-(4-toluenesulfonyl)-4-(toluenesulfonyloxy)-D-proline ethyl ester (154.0 g, 0.33 mole) and anhydrous tetramethylammonium acetate (56.15 g, 0.42 mole) in toluene (1000 mL) was refluxed under nitrogen atmosphere for 18 h. The cooled reaction mixture was extracted with H₂O (2 x 300 mL). The organic phase was dried over MgSO₄, filtered, and concentrated under reduced pressure. The residue was treated with i-PrOH (200 mL) and cooled to 0 °C for 30 min. The resulting crystalline product was collected by filtration to give the title compound (86.78 g, 52%).

(5) 4-Hydroxy-1-(4-toluenesulfonyl)-D-proline ethyl ester. To a suspension of 4-(acetyloxy)-1-(4-toluenesulfonyl)-D-proline ethyl ester (86.0 g, 0.24 mole) in MeOH (1000 mL), was added water (415 mL). The pH was adjusted to 11 by the addition of Na₂CO₃ (6g). After 4 h the pH was adjusted to 7 by the addition of AcOH (1.5 mL) and the mixture was kept at 23 °C for 17 h. The pH was adjusted to 7 with AcOH (0.5 mL) and the volume of the solution was reduced by 50% by evaporation under reduced pressure. The solution was extracted with CH₂Cl₂ (3 x 750 mL). The combined organic extracts were washed with H₂O, dried over MgSO₄, filtered, and evaporated under reduced pressure to give a mixture of 4-Hydroxy-1-(4-toluenesulfonyl)-D-proline ethyl and methyl esters (68.16 g) as a viscous oil.

45 (6) (2R,4S)-1-(4-toluenesulfonyl)-2-hydroxymethyl-4-hydroxy pyrrolidine. To a cold solution of 4-hydroxy-1-(4-toluenesulfonyl)-D-proline ethyl and methyl ester (68.0 g, 0.22 mole) in THF (650 mL) was added LiBH₄ (16.7 g, 0.77 mole) in portions. The reaction mixture was stirred at 0 °C for 1 h and allowed to warm to 23 °C and stand for 20 h. The mixture was cooled to 0 °C and the pH was adjusted to 3 with 6N HCl. The volume was reduced to 300 mL in vacuo and the white precipitate that formed was filtered,
50 washed with cold water, and dried in vacuo at 50 °C for 24 h yielding the title compound (58.5 g, 99.4%) which was used in the next step without further purification.

(7) (2R,4S)-1-(4-toluenesulfonyl)-2-(4-toluene-sulfonyloxymethyl)-4-(4-toluenesulfonyloxy)-pyrrolidine. p-Toluenesulfonyl chloride (142.99 g, 0.75 mole) was added portionwise to a solution of (2R,4S)-1-(4-toluenesulfonyl) -2-hydroxymethyl-4-hydroxy pyrrolidine (58.0 g, 0.21 mole) in pyridine (350 mL) at 0 °C. The reaction mixture was kept at 5 °C for 2 h and was stirred at 23 °C for 20 h. The mixture was poured into 2.4 N HCl (1500 mL) and extracted with CH₂Cl₂ (3 x 750 mL). The combined CH₂Cl₂ extracts were dried K₂CO₃, filtered, and concentrated under reduced pressure. Trituration of the residue with EtOH afforded a solid which was collected by filtration, and dried in vacuo at 65 °C for 24 h to give the title

55

compound (74.17 g, 60.9% m.p. 130°-132° C).

(8) (1R,4R)-2-(4-toluenesulfonyl)-5-phenylmethyl-2, 5-diazabicyclo[2.2.1]heptane. A mixture of (2R,4S)-1-(4-toluenesulfonyl)-2-(4-toluenesulfonyloxymethyl)-4-(4-toluenesulfonyloxy)-pyrrolidine (73.5 g, 0.13 mole) and benzylamine (44.84 g, 0.42 mole) in toluene (350 mL) was heated at reflux for 20 h while under a N₂ atmosphere. The reaction mixture was cooled to 23° C, filtered and the collected precipitate was washed with toluene (250 mL). The combined organic filtrates were concentrated under reduced pressure. The residue was dissolved in i-PrOH (50 mL) and the solid that formed was collected by filtration to give the title compound (38.95 g, 87.6%, [α]_D = -16.82 (c = 1, acetone)).

(9) (1R,4R)-5-phenylmethyl-2,5-diazabicyclo[2.2.1] heptane dihydrobromide. A mixture of (1R,4R)-2-(4-toluenesulfonyl)-5-phenylmethyl-2,5-diazabicyclo[2.2.1]-heptane (38.95 g, 0.11 mole) in AcOH (500 mL) containing HBr (30% wt) was heated at 70° C for 18 h. The mixture was allowed to cool and the precipitate that formed was filtered and washed with acetone. The filtrate was concentrated to one-third of the original volume and the resulting solid was combined with the first precipitate to give the title compound (36.55 g, 92%, m.p. 272°-275° C).

15

Anal. Calcd. for C ₁₂ H ₁₆ N ₂ ·2HBr:	
	C, 41.17; H, 5.19; N, 8.01.
Found:	C, 41.05; H, 5.16; N, 7.76.

20

(10) 2-(tert-butyloxycarbonyl)-5-phenylmethyl-(1R,4R)-2,5-diazabicyclo[2.2.1]heptane. Into a solution of (1R,4R)-5-phenylmethyl-2,5-diazabicyclo[2.2.1]heptane (3.0 g, 15.96 mmol) in CH₂Cl₂ (15 mL) at 0° C was added dropwise di-tert-butylidicarbonate (3.83 g, 17.55 mmol) in 10 mL of CH₂Cl₂. After the addition was complete, the reaction mixture was stirred at 23° C for 19 h. The reaction mixture was washed with saturated NaHCO₃ (3 x 25 mL) followed by saturated NaCl solution, dried over K₂CO₃, filtered and concentrated under reduced pressure to give the title compound (4.62 g). The product was used in the next step without further purification.

(11) 2-(t-butyloxycarbonyl)-(1R,4R)-2,5-diazabicyclo [2.2.1]heptane. Into a solution of (10) (3.65 g, 12.67 mmol) in 200 mL of EtOH was added AcOH (1 mL). The reaction mixture was treated with 10% palladium-on-carbon (1.10 g) and hydrogenated at 50 psi for 2 h at 23° C. The mixture was then heated to 60° C and hydrogenation was continued for 2 h. After this time, heating was discontinued; the reaction mixture was allowed to cool to room temperature and hydrogenation was continued for 3 h. The reaction mixture was filtered through celite and the filtrate was concentrated under reduced pressure. The residue was made alkaline by the addition of 5N NaOH and extracted with CH₂Cl₂ (5 x 100 mL). Combined CH₂Cl₂ extracts were dried over K₂CO₃, filtered and concentrated under reduced pressure to give the title compound (2.34 g, 93.2%). The product was used in the next step without further purification.

(12) 2-(t-butyloxycarbonyl)-5-(5-fluoro-4-methylthio-2-pyrimidinyl)-(1R,4R)-2,5-diazabicyclo[2.2.1]-heptane. The title compound was prepared as described for the 1S,4S isomer Example 7 (10). Yield: 6.80 g, 88.2%.

(13) 2-(5-fluoro-2-pyrimidinyl)-(1R,4R)-2,5-diazabicyclo[2.2.1]heptane. The title compound was prepared as described for the 1S,4S isomer Example 7 (11). Yield: 2.74 g, 88.4%.

Other intermediates and starting materials used for preparation of Formula I compounds are either available commercially or are readily available to one skilled in the art via the chemical literature or by appropriate modification of the foregoing examples.

SYNTHESIS OF PRODUCTS

50

Formula IA Compounds

55

EXAMPLE 9

1-(4-(4-Fluorophenyl)butyl)-4-(5-fluoro-2-pyrimidinyl) piperazine

A mixture of 4-(5-fluoro-2-pyrimidinyl)piperazine (1.66 g, 9.1 mmol), 1-chloro-4-(4-fluorophenyl)butane (1.68 g, 9.0 mmol), K_2CO_3 (2.48 g, 17.9 mmol), KI (0.15 g, 0.9 mmol) and MeCN (90 mL) was refluxed under N_2 atm. for 40 h and then allowed to cool at 23 °C and filtered; the solids were washed with MeCN (2 x 20 mL). The filtrates were combined and concentrated under reduced pressure to a gum which was dissolved in EtOAc (200 mL). The organic solution was washed with water (20 mL), dried ($MgSO_4$) and concentrated under reduced pressure. The residue was passed through a silica gel pad using EtOAc acetate as eluting solvent. The appropriate fractions were combined and concentrated; the resulting solid was recrystallized from ethanol to afford analytical sample; mp 59-61 °C; 1H NMR ($CDCl_3$, 200 MHz) δ : 1.4-1.8 (m, 4H), 2.2-2.5 (m, 6H), 2.45 (t, J=5.1 Hz), 2.5-2.7 (m, 2H), 3.74 (t, J=5.1 Hz, 4H), 6.9-7.0 (m, 2H), 7.0-7.2 (m, 2H), 8.17 (s, 2H); IR (KBr) ν : 1610, 1555, 1511, 1485, 1455, 1445, 1360 cm^{-1} ;

Anal. Calcd. for $C_{18}H_{22}N_4F_2$:	
Found:	C, 65.04; H, 6.67; N, 16.86. C, 64.95; H, 6.77; N, 16.85.

EXAMPLE 10

1-[3-(4-Fluorophenylthio)propyl]-4-(5-fluoro-2-pyrimidinyl) piperazine

(1) 3-(4-fluorophenylthio)-1-propanol. A mixture of 4-fluorophenylthiol (15 g, 0.117 mole), 3-chloro-1-propanol (10.8 mL, 0.128 mole) and N^2OH (4.96 g, 0.124 mole) in EtOH (120 mL) was refluxed under N_2 atm. for 20 h, cooled to 23 °C and filtered. The insoluble material was washed with EtOH (10 mL). The filtrate and washings were concentrated under reduced pressure to a crude material (23.3 g) which was distilled under high vacuum, 17.0 g (78%), bp 120-2 °C/0.75 mmHg; 1H NMR ($CDCl_3$, 200 MHz) δ : 1.44 (t, J=5.3 Hz, 1H), 1.75-1.0 (m, 2H), 2.97 (t, J=7.1 Hz, 2H), 3.72 (m, 2H), 6.9-7.1 (m, 2H), 7.3-7.5 (m, 2H); IR (film) ν : 3600-3000, 1590, 1490, 1225 cm^{-1} ;

Anal. Calcd. for $C_9H_9 \cdot OFS \cdot 0.1H_2O$:	
Found:	C, 57.48; H, 6.00; S, 17.05. C, 57.47; H, 5.89; S, 16.73.

(2) 1-bromo-3-(4-fluorophenylthio)propane. A mixture of 3-(4-fluorophenylthio)-1-propanol (9.1 g, 48.9 mmol), aqueous HBr (48%, 14 mL) and conc. aqueous H_2SO_4 (2.4 mL) was refluxed for 24 h, cooled and carefully poured onto ice-water mixture (120 mL). The aqueous phase was extracted with Et_2O (3 x 25 mL). The Et_2O layer was dried ($MgSO_4$) and concentrated to a crude oil which was distilled under high vacuum, 6.9 g, (57%), bp 108 °-114 °C/0.6 mmHg; 1H NMR ($CDCl_3$, 200 MHz) δ : 2.0-2.2 (m, 2H), 3.00 (t, J=6.9 Hz, 2H), 3.50 (t, J=6.3 Hz, 2H), 6.9-7.1 (m, 2H), 7.3-7.5 (m, 2H); IR (film) ν : 1590, 1490, 1225 cm^{-1} ;

Anal. Calcd. for C_9H_9BrFS :	
Found:	C, 43.39; H, 4.05. C, 41.31; H, 3.88.

(3) A mixture of 4-(5-fluoro-2-pyrimidinyl)piperazine (1.58 g, 8.67 mmol), 1-bromo-3-(4-fluorophenylthio)propane (2.16 g, 8.67 mmol), triethylamine (1.33 mL, 9.53 mmol) and KI (1.58 g, 9.52 mmol) in MeCN (45 mL) was refluxed under N₂ atm. for 18 h, then cooled at 23 °C and concentrated under reduced pressure. The residue was solubilized in EtOAc (400 mL) and the resulting organic solution was washed with water (2 x 40 mL), dried (MgSO₄) and concentrated to dryness. The crude material (3.0 g) was purified on a silica gel pad (3.4 x 8.5 cm) using a mixture of 50%-100% EtOAc in hexane. Appropriate fractions were combined and concentrated leaving 2.64 g (87%) mp 68 °-70 °C. Recrystallization from EtOH afforded an analytical sample, mp 70-1 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 1.6-1.9 (m, 2H), 2.2-2.7 (m, 6H), 2.92 (t, J=7 Hz, 2H), 3.6-3.9 (m, 4H), 6.9-7.1 (m, 2H), 7.3-7.5 (m, 2H), 8.17 (s, 2H); IR (KBr) ν: 1609, 1552, 1500, 1490 cm⁻¹;

Anal. Calcd. for C ₁₇ H ₂₉ N ₄ SF ₂ :	
Found:	C, 58.27; H, 5.75; N, 15.99; S, 9.15. C, 58.21; H, 5.73; N, 15.87; S, 9.43.

EXAMPLE 11

1-(3-(1,3-dioxolan-2-yl)propyl)-4-(5-Fluoro-2-pyrimidinyl) piperazine.

A mixture of 4-(5-fluoro-2-pyrimidinyl)piperazine (7.29 g, 40.0 mmol), 2-(3-chloropropyl)-1,3-dioxolane (6.62 g, 5.8 mL, 44.0 mmol) and triethylamine (12.5 mL, 90.0 mmol) in 2-butanone (250 mL) was heated to reflux and treated dropwise (1 h) with a solution of NaI (3.0 g, 20.0 mmol) in 2-butanone (70 mL). The reaction mixture was refluxed for 12 h then treated dropwise (1 h) with a solution of NaI (3.0 g, 20.0 mmol) in 2-butanone (70 mL) and refluxed for another 12 h. After cooling at 23 °C, the solvent was evaporated in vacuo leaving a residue which was solubilized in EtOAc (400 mL). The organic solution was washed with aqueous NaOH solution (20 mL, 2N); the aqueous phase was extracted with EtOAc (2 x 50 mL). The organic extracts were combined, dried (MgSO₄) and concentrated to dryness. The solution was chromatographed on a silica gel column (4.5 x 15 cm) using a mixture of 20% MeCN in EtOAc. Appropriate fractions were concentrated in vacuo leaving a pale yellow syrup which crystallized on standing, 10.5 g, mp 54-5 °C (89%). Recrystallization from EtOAc-pet.ether mixture (1:14) gave an analytical sample, mp 55-6 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 1.5-1.9 (m, 4H), 2.3-2.7 (m, 6H), 3.6-4.1 (m, 8H), 4.8-5.0 (m, 1H), 8.16 (s, 2H); IR(KBr) ν: 1610, 1555, 1489 cm⁻¹; UV (EtOH) λ: 244 (ε 17239), 332 (ε 1948);

Anal. Calcd. for C ₁₄ H ₂₁ N ₄ O ₂ F:	
Found:	C, 56.74; H, 7.14; N, 18.91. C, 56.77; H, 7.15; N, 18.98.

EXAMPLE 12

1-(2-Thienyl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl] butanone

To a hot (78 °C) mixture of 4-(5-fluoro-2-pyrimidinyl) piperazine (4.00 g, 22.0 mmol), 4-chloro-2'-butyrothienone (3.57 mL, 22.0 mmol) and triethylamine (6.97 mL, 50.0 mmol) in methylethyl ketone (125 mL) was added dropwise (1 h) a solution of NaI (4.8 g, 32.0 mmol) in methylethyl ketone (125 mL). The resulting mixture was refluxed for 22 h, then cooled at 23 °C and concentrated to dryness in vacuo. The

residue was diluted with CH_2Cl_2 (400 mL) and the solution washed with aqueous NaOH solution (0.5 N, 44 mL). Aqueous phase was separated and extracted with CH_2Cl_2 (2 x 10 mL). Organic extracts were combined, dried (MgSO_4) and concentrated under reduced pressure to a crude material which was chromatographed on a silica gel column (5 x 16 cm) using a mixture 30-100% EtOAc in hexane. Appropriate fractions were combined and concentrated to afford a yellow solid, 2.0 g (27%). Recrystallization from Et_2O gave an analytical sample, mp 77-8° C. ^1H NMR (CDCl_3 , 200 MHz) δ : 1.9-2.2 (m, $\text{CH}_2\text{CH}_2\text{CH}_2$, 2H), 2.3-2.7 (m, 6H), 2.96 (t, $J = 7.0$ Hz, COCH_2 , 2H), 3.6-3.9 (m, 4H), 7.0-7.2 (m, 1H), 7.5-7.7 (m, 1H), 7.7-7.8 (m, 1H), 8.18 (s, pyrimidinyl H, 2H); IR (KBr) ν : 660, 1610, 1555, 1510, 1480, 1359 cm^{-1} ; UV (EtOH) λ : 248 (ϵ 19699), 284 (ϵ 6287), 330 (ϵ 1816).

Anal. Calcd. for $\text{C}_{15}\text{H}_{15}\text{N}_4\text{OSF}$:	
	C, 57.47; H, 5.73; N, 16.75; S, 9.59.
Found:	C, 57.42; H, 5.75; N, 16.81; S, 9.77.

EXAMPLE 13

1-(4-Fluoronaphth-1-yl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl]butanone

(1) Ethyl 4-(4-(5-fluoropyrimidin-2-yl)piperazin-1-yl) butanoate

A mixture of 4-(5-fluoro-2-pyrimidinyl)piperazine (1.37 g, 7.5 mmol), ethyl 4-bromobutanoate (1.07 mL, 7.5 mmol), triethylamine (1.35 mL, 9.7 mmol) in 2-butanone (50 mL) was refluxed under Ar atm. for 4 h. After cooling at 23° C, the reaction mixture was filtered and the filtrate concentrated under reduced pressure to a crude mixture which was dissolved in EtOAc (150 mL). The organic solution was washed with water (2 x 15 mL), dried (MgSO_4) and concentrated under reduced pressure to a crude mixture which was purified on a silica gel pad using a mixture 0-100% EtOAc in hexane as eluting solvent. Evaporation of appropriate fractions gave 1.66 g (75%) of pure compound; ^1H NMR (CDCl_3 , 200 MHz) δ : 1.25 (t, $J = 7.1$ Hz, CH_3 , 3H), 1.84 (t of t, $J = 7.1$ Hz, H-3, 2H), 2.2-2.61 (m, 8H), 3.6-3.9 (m, 4H), 4.13 (q, $J = 7.1$ Hz, CH_2CH_3 , 2H), 8.17 (s, aromatic H, 2H); IR (film) ν : 1732 (C=O), 1610, 1552, 1500, 1360 cm^{-1} ;

Anal. Calcd. for $\text{C}_{14}\text{H}_{22}\text{N}_4\text{O}_2\text{F}$:	
	C, 56.74; H, 7.14; N, 18.91.
Found:	C, 56.63; H, 7.27; N, 18.66.

(2) Ethyl 2-(4-fluorobenzoyl)-4-(4-(5-fluoro-2-pyrimidinyl)piperazin-1-yl)butanoate

To a cold (-78° C) solution of lithium 1,1,1,3,3,3-hexamethyl disilazane in tetrahydrofuran (27.4 mL, 1N, 27.4 mmol) kept under Ar atm was added dropwise (10 min) a solution of ethyl 4-(4-(5-fluoro-2-pyrimidinyl)piperazin-1-yl)butanoate (3.7 g, 12.5 mmol) in dry THF (12 mL). The reaction mixture was stirred 0.25 h at -78° C and then treated dropwise (10 min) with a solution of 4-fluoro-1-naphthoyl chloride (2.6 g, 12.5 mmol) in dry THF (10 mL). The reaction was stirred -78° C for 0.5 h then the cooling bath was removed. When the temperature of reaction mixture reached 0° C, HCl solution (40 mL, 0.3N) was added slowly followed by CH_2Cl_2 (350 mL); the organic phase was separated and aqueous phase was extracted with CH_2Cl_2 (120 mL). The organic extracts were combined, washed with water (2 x 45 mL) and brine, dried (MgSO_4) and concentrated under reduced pressure to a crude mixture which was purified on a silica gel pad (8.7 x 2.5 cm). Evaporation of appropriate fractions gave 2.8 g (48%) of the desired product compound which was

recrystallized from EtOH to give an analytical sample; mp 95-6 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 1.08 (t, J=7.1 Hz, CH₃, 3H), 2.1-2.7 (m, 8H), 3.5-3.8 (m, 4H), 3.9-4.3 (m, 2H), 4.5-4.7 (m, 1H), 7.1-7.25 (m, 1H), 7.5-7.8 (m, 2H), 8.17 (s), 8.0-8.02 (s and m, 4H), 8.6-8.7 (m, 1H); IR (KBr) ν: 1740 (C=O), 1682, 1630, 1611, 1600, 1578, 1554, 1510 cm⁻¹; UV (CH₃CN) λ: 224 (ε 43570), 242 (ε 37740), 304 (ε 9084);

Anal. Calcd. for C ₂₅ H ₂₆ N ₄ O ₃ F ₂ :	
	C, 64.09; H, 5.59; N, 11.96.
Found:	C, 63.98; H, 5.62; N, 11.90.

and 1.74 g (51%) of the by-product coupled compound: 1,7-bis [4-(5-fluoropyrimidin-2-yl)piperazin-1-yl]-3-ethoxycarbonylheptan-4-one. Treatment of an ethanolic solution of the oil with two equivalents of ethanolic HCl afforded the bis hydrochloride salt; mp 196-8 °C;

(3) A mixture of Ethyl 2-(4-fluorobenzoyl)-4-(4-(5-fluoro-2-pyrimidinyl)piperazin-1-yl)butanoate (2.95 g, 6.3 mmol) in aqueous HCl (65 mL, 1N) was refluxed for 2 h, cooled at 23 °C and basified, first with aqueous NaOH solution (30 mL, 2N) and then with saturated NaHCO₃ solution until pH=9. The aqueous solution was extracted with CH₂Cl₂ (3 x 150 mL). Organic extracts were combined, washed with water (30 mL) and brine, dried (MgSO₄) and concentrated to dryness. The crude material was chromatographed on silica gel using a mixture of 50% CH₂Cl₂ in EtOAc. Appropriate fractions were combined and solvent evaporated under reduced pressure leaving 2.3 g (92%), mp 119-120 °C.

Recrystallization from EtOH afforded analytical sample; mp 120-1 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 2.03 (t, J=7.0 Hz, CH₂CH₂CH₂, 2H), 2.4-2.6 (m, 6H), 3H (t, J=7.0, CH₂CO, 2H), 3.8-3.9 (m, 4H), 7.1-7.2 (m, 1H), 7.5-7.8 (m, 2H), 7.9-8.0 (m, 1H), 8.18 (s), 8.1-8.3 (m, 3H), 8.7-8.8 (m, 1H); IR (KBr) ν: 1681, 1630, 1611, 1600, 1575, 1555, 1494 cm⁻¹; UV (CH₃CN) λ: 226 (ε 40377), 298 (ε 7498);

Anal. Calcd. for C ₂₂ H ₂₂ N ₄ OF ₂ :	
	C, 66.65; H, 5.59; N, 14.13.
Found:	C, 66.47; H, 5.59; N, 14.08.

EXAMPLE 14

4-[5-(5-Fluoro-2-pyrimidinyl)-(1S,4S)-2,5-diazabicyclo[2.2.1] heptan-2-yl]-1-(4-fluorophenyl)butanone

A mixture of 2-(5-fluoro-2-pyrimidinyl)-(1S,4S)-2,5-diazabicyclo[2.2.1]heptane (Ex. 7(11)), 4-chloro-4'-fluorobutyrophenone ethylene ketal (3.53 g, 14.43 mmol), micropulverized K₂CO₃ (5.97 g, 43.29 mmol), and KI (0.36 g, 2.16 mmol) in 50 mL of MeCN was refluxed for 22 h. K₂CO₃ was filtered and the reaction mixture was concentrated in vacuo. 3N HCl (7 mL) was added to the residue. The reaction mixture was heated on a steam bath for 15 min. Then, 20 mL of EtOH was added and heating was continued for 30 min. The solution was concentrated and the residue was triturated with EtOH/Hexane which induced crystallization. The solid was collected by suction filtration to give the hydrochloride salt of the title compound (2.96 g, mp 192-196 °C dec). The salt was converted to its free base to give the title compound (2.15 g, 41.6%).

EXAMPLE 15

4-[5-(5-Fluoro-2-pyrimidinyl)-(1R,4R)-2,5-diazabicyclo [2.2.1]heptan-2-yl]-1-(4-fluorophenyl)butanone

The title compound was prepared as described for the 1S,4S isomer (Ex. 14). Yield: 2.40 g, 49.3%.

EXAMPLE 16

5 4-[5-(5-Fluoro-2-pyrimidinyl)-(1S,4S)-2,5-diazabicyclo [2.2.1]heptan-2-yl]-1-(4-fluorophenyl)butanol hydro-
chloride

Into a solution of the ketone (Ex. 14) (2.15 g, 6.01 mmol) in 140 mL of EtOH was added NaBH₄ (0.68 g, 18.02 mmol). The reaction mixture was stirred at 23 °C for 2 h and refluxed for 0.5 h and concentrated in
 10 vacuo. The residue was taken up in CH₂Cl₂ (50 mL) and H₂O (20 mL). The aqueous layer was made basic
 with 30% NaOH solution and extracted with CH₂Cl₂ (3 x 100 mL). Combined CH₂Cl₂ extracts were dried
 over K₂CO₃, filtered and concentrated under reduced pressure. Flash chromatography (CH₂Cl₂: MeOH;
 93:7) gave the title compound (1.18 g, 54.6%). Conversion to the hydrochloride salt followed by recrystal-
 lization from EtOH-Et₂O afforded a white solid, mp 186-191 °C.

15

Anal. Calc. for C ₉ H ₂₂ F ₂ N ₄ O.HCl:	
	C, 57.51; H, 5.85; N, 14.12.
Found:	C, 57.35; H, 5.89; N, 14.10.

20

EXAMPLE 17

25

30 4-[5-(5-Fluoro-2-pyrimidinyl)-(1R,4R)-2,5-diazabicyclo [2.2.1]heptan-2-yl]-1-(4-fluorophenyl)butanol hydro-
chloride

The title compound was prepared as described for the 1S,4S isomer (Ex. 16). Yield: 1.17 g (48.5%), mp
 187-191 °C.

35

Anal. Calc. for C ₉ H ₂₂ F ₂ N ₄ O.HCl:	
	C, 57.51; H, 5.85; N, 14.12.
Found:	C, 57.46; H, 5.89; N, 14.21.

40

Formula 1B Compounds

45

EXAMPLE 18

50 1-[4-Acetamido-4-(4-fluorophenyl)butyl]-4-(5-fluoro-2-pyrimidinyl)piperazine monohydrochloride

(1) 1-[4-azido-4-(4-fluorophenyl)butyl]-4-(5-fluoro-2-pyrimidinyl)piperazine monohydrochloride

55 To a cold (5 °C) solution of triphenylphosphine (12.6 g, 48.0 mmol) in dry THF (130 mL) kept under an
 Ar atmosphere was added dropwise over 1 h period diisopropyl azodicarboxylate (9.5 mL, 48.2 mmol)
 followed by a solution of 1-(4-fluorophenyl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl] butanol-1-01 Ex. 5;
 (14.6 g, 41.9 mmol) in dry THF (275 mL). The reaction mixture was immediately treated dropwise (10 min)

with a solution of diphenylphosphoryl azide (13.3 g, 48.3 mmol) in dry THF (130 mL). The resulting reaction mixture was stirred at 5 °C for 1 h then at 23 °C for 16 h before being concentrated to dryness. The residue was suspended in CH₂Cl₂ (200 mL) and filtered. The filtrate was concentrated and chromatographed on a silica gel pad using a mixture 30-100% EtOAc in CH₂Cl₂ as eluting solvent. Appropriate fractions were concentrated under reduced pressure leaving 8.0 g, 51%; ¹H NMR (CDCl₃, 200 MHz) δ: 1.3-2.0 (m, 4H), 2.2-2.6 (m, 6H), 3.6-3.9 (m, 4H), 4.43 (t, J = y.1 Hz, 1H), 6.9-7.1 (m, 2H), 7.1-7.4 (m, 2H), 8.16 (s, 2H). A part of the compound was converted to monohydrochloride salt and recrystallized from EtOH-Et₂O mixture to afford an analytical sample; mp 176-8 °C; ¹H NMR (DMSO-d₆, 200 MHz) δ: 1.5-2.0 (m, 4H), 2.8-3.2 (m, 4H), 3.2-3.6 (m, 4H), 4.4-4.7 (m, 2H), 4.7-4.9 (m, 1H), 7.2-7.4 (m, 2H), 7.4-7.6 (m, 2H), 8.5 (s, 2H), 10.5-10.9 (m, 1H); IR (KBr) ν: 2810-2300, 2100, 1605, 1560, 1510, 1475, 1440 cm⁻¹; UV (H₂O) λ: 238 (ε 16600), 320 (ε 2011);

Anal. Calcd. for C ₁₈ H ₂₁ N ₇ F ₂ .HCl.0.15 H ₂ O:	
---	--

Found:	C, 52.40; H, 5.45; N, 23.76. C, 52.58; H, 5.34; N, 23.88.
--------	--

(2) 1-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazinybutanamine

A mixture of 1-[4-azido-4-(4-fluorophenyl)butyl]-4-(5-fluoro-2-pyrimidinyl)piperazine (3.3 g, 8.84 mmol), 10% palladium on charcoal (0.33 g) in EtOH (70 mL) was hydrogenated under 40 psi at 23 °C for 2h. The mixture was filtered and solvent concentrated in vacuo leaving the title compound, 2.93 g (95%); IR (film) ν: 1610, 1552, 1505 cm⁻¹.

(3) To a cold (5 °C) solution of 1-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazinybutanamine (2.33 g, 6.7 mmol) in glacial AcOH (10 mL) was added dropwise (20 min) acetic anhydride (0.64 mL, 6.8 mmol). The cooling bath was removed and the reaction mixture was stirred at 23 °C for 16 h, concentrated to half of its initial volume and diluted in CH₂Cl₂ (100 mL). The organic solution was basified with the addition of saturated NaHCO₃ aqueous solution until pH 9. The organic layer was separated and the aqueous layer was extracted with CH₂Cl₂ (2 x 50 mL). Organic layers were washed with brine, dried (MgSO₄) and concentrated to dryness. The resulting residue was chromatographed on a silica gel pad (8.7 x 4.5 m) using a mixture of 5-20% EtOH in CHCl₃. Appropriate fractions were concentrated in vacuo leaving a solid, 2.0 g (77%). Recrystallization from EtOH-Et₂O mixture gave an analytical sample, mp 142-3 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 1.3-1.7 (m, 2H), 1.7-1.9 (m, 2H), 1.99 (s, NHCOCH₃, 3H), 2.3-2.6 (m, 6H), 3.7-3.9 (m, 4H), 4.9-5.1 (m, CHNH, 1H), 6.14 (bd, J = 6.7 Hz CHNH, 1H), 6.9-7.1 (m, 2H), 7.1-7.3 (m, 2H), 8.18 (s, pyrimidinyl H, 2H); IR (KBr) ν: 3290, 1650, 1610, 1551, 1510, 1500, 1395, 1356 cm⁻¹; UV (CH₃CN) λ: 246 (ε 19523), 332 (ε 1981);

Anal. Calcd. for C ₂₀ H ₂₅ NOF ₂ :	
---	--

Found:	C, 61.68; H, 6.47; N, 17.98. C, 61.82; H, 6.54; N, 18.02.
--------	--

EXAMPLE 19

1-[4-(4-Fluorobenzamido)-4-(4-fluorophenyl)butyl]-4(5-fluoro-2-pyrimidinyl)piperazine

To a cold (5 °C) solution of 1-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazinybutanamine (1.93 g, 5.55 mmol) in CH₂Cl₂ (40 mL) was added triethylamine (0.78 mL, 5.6 mmol) and 4-fluorobenzoyl chloride

(0.66 mL, 5.6 mmol) over 30 min period. The reaction mixture was stirred at 5 °C for 1 h then at 23 °C for 1 h before being diluted with CH₂Cl₂ chloride (100 mL). The resulting organic solution was washed with cold (0 °C) water (20 mL), aqueous NaHCO₃ saturated solution (pH 9) and brine, dried (MgSO₄) and concentrated to dryness. The crude mixture was chromatographed on a silica gel pad using a mixture of 4-10% EtOH in CH₂Cl₂. Appropriate fractions were concentrated to afford a white solid, 2.0 g (77%). Analytical sample was prepared from recrystallization from EtOH, mp 147-8 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 1.4-1.8 (m, 2H), 1.8-2.2 (m, 2H), 2.3-2.6 (m, 6H), 3.6-3.9 (m, 4H), 5.0-5.3 (m, CH₂CHNH, 1H), 6.6 (d, J=7.3 Hz, CHNH, 1H), 6.9-7.2 (m, 4H), 7.26-7.4 (m, 2H), 7.7-7.9 (m, 2H), 8.18 (s, pyrimidinyl H, 2H); IR (MBr) ν: 3330, 1632, 1605, 1552, 1505, 1395, 1358 cm⁻¹; UV (CH₃CN) λ:244 (ε 25182), 332 (ε 1922);

Anal. Calcd. for C ₂₅ H ₂₆ N ₅ OF ₃ :	
	C, 63.96; H, 5.58; N, 14.92.
Found:	C, 64.07; H, 5.59; N, 14.90.

EXAMPLE 20

1-[4-(4-Fluorophenyl)-4-(4-fluorophenylsulfonamido) butyl]-4-(5-fluoro-2-pyrimidinyl)piperazine

To a cold (5 °C) solution of 1-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazinybutanamine (2.34 g, 6.73 mmol) in CH₂Cl₂ (30 mL) was added triethylamine (0.94 mL, 6.7 mmol) and a solution of 4-fluorobenzene sulfonyl chloride (1.31 g, 6.73 mmol) in CH₂Cl₂ (15 mL) over 0.5 h. The reaction mixture was stirred at 5 °C for 1 h then at 25 °C for 1 h before being diluted with CH₂Cl₂ (160 mL). The resulting organic solution was washed with cold water (0 °C) (2 x 20 mL), aqueous NaHCO₃ saturated solution (pH 9) and brine, dried (MgSO₄) and concentrated to dryness. The crude material was chromatographed on a silica gel pad using a mixture of 5-10% EtOH in CH₂Cl₂. The appropriate fractions were concentrated in vacuo leaving a white solid, 2.8 g (82%). Recrystallization from EtOH afforded an analytical sample, mp 128-9 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 1.3-1.7 (m, 2H), 1.7-1.9 (m, 1H), 1.9-2.1 (m, 1H), 2.2-2.8 (m, 6H), 3.6-4.2 (m, 4H), 4.48 (bs, 1H), 6.8-7.2 (m, 6H), 7.5-7.8 (m, 2H), 8.20 (s, pyrimidinyl H, 2H), 8.58 (bs, NH, 1H); IR (KBr) ν: 3270, 1610, 1592, 1554, 1509, 1492, 1439, 1400 cm⁻¹; UV (CH₃CN) λ:244 (ε 19445), 330 (ε 1915);

Anal. Calcd. for C ₂₄ H ₂₆ N ₅ O ₂ F ₃ S:	
	C, 57.02; H, 5.18; N, 13.85; S, 6.34.
Found:	C, 56.89; H, 5.23; N, 13.87; S, 6.58.

EXAMPLE 21

1-[3-(4-Fluorophenylsulfonyl)propyl]-4-(5-fluoro-2-pyrimidinyl)piperazine

To a cold (5 °C) solution of 1-[3-(4-fluorophenylthio) propyl]-4-(5-fluoro-2-pyrimidinyl)piperazine (3.55 g, 10.1 mmol) in acetic acid (6 mL) was added ammonium molybdate (0.08 g, 0.4 mmol) and 30% H₂O₂ solution (2.8 mL, 27.4 mmol) was added dropwise at such a rate that the temperature was kept at 10 °C (~ 3 h). The reaction mixture was then stirred at 23 °C for 16 h before being diluted with water (120 mL) and CH₂Cl₂ (200 mL). Excess H₂O₂ was destroyed by addition of aqueous saturated sodium sulfite solution. The mixture was basified with Na₂CO₃ (pH 8.5). The organic phase was separated and the aqueous phase further extracted with CH₂Cl₂ (2 x 100 mL). The organic extracts were combined, washed with brine, dried

(MgSO₄) and concentrated under reduced pressure leaving a crude mixture which was purified on a silica gel pad using a mixture of 2% EtOH in CHCl₃ as eluting solvent. The appropriate fractions were concentrated in *vacuo* leaving a white solid 3.6 g, 93% which was recrystallized from EtOH given an analytical sample; mp 132.5-3.5 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 1.8-2.0 (m, 2H), 2.3-2.6 (m, 6H) 3.1-3.3 (m, 2H), 3.6-3.8 (m, 4H), 7.1-7.3 (m, 2H), 7.8-8.0 (m, 2H), 8.16 (s, 2H); IR (KBr) ν: 1612, 1590, 1558, 1491, 1470, 1450, 1358, (SO₂), 1145 (SO₂) cm⁻¹; UV (CH₃CN) λ: 218 (ε 12429), 246 (ε 17791), 330.

EXAMPLE 22

1-Cyclohexyl-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl] butanol

(1) 4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl] butyraldehyde

A mixture of 4-(4-(5-fluoro-2-pyrimidinyl)piperazin-1-yl) butyraldehyde ethylen acetal (1.36 g, 4.6 mmol) and aqueous HCl (6N, 8 mL, 48.0 mmol) was stirred at 45 °C for 2 h and then basified after cooling with aqueous NaOH solution (2N, 22-24 mL, ~pH 10). The resulting mixture was extracted with CH₂Cl₂ (3 x 50 mL) leaving 1.3 g of crude oil after evaporation of solvent. Purification on a silica gel column (2.5 x 10 cm) using a mixture of 20% MeCN in EtOAc afforded 0.86 g (74%) of pale yellow syrup; ¹H NMR (CDCl₃, 200 MHz) δ: 1.86 (m, 2H), 2.39 (t, J = 6.9 Hz), 2.3-2.6 (t, m, 8H), 3.74 (m, 4H), 8.18 (s, 2H), 9.8 (t, J = 1.6 Hz, 1H).

(2) To a cold (-30 °C) solution of 4-(4-(5-fluoropyrimidinyl)-piperazin-1-yl) butyraldehyde (4.32 g, 17.1 mmol) in THF kept under N₂ atmosphere was added dropwise (0.5 h) a solution of cyclohexyl magnesium chloride in Et₂O (17 mL, 2N, 34.0 mmol). The reaction mixture was stirred at -30 °C for 1 h before adding water dropwise (35 mL) and EtOAc (350 mL). The organic phase was separated and the aqueous phase extracted with EtOAc (200 mL). The organic extracts were dried (MgSO₄) and concentrated under reduced pressure to a yellow syrup (6.1 g). Purification on silica gel column (4.5 x 17 cm) gave 2.54 g (44%) of a colorless syrup which crystallized (mp 80-2 °C) on standing. Recrystallization from EtOAc - pet. ether mixture afforded an analytical sample; mp 81-3 °C; ¹H NMR (CDCl₃, 20 MHz) δ: 0.8-1.5 (m, 8H), 1.5-2.0 (m, 7H), 2.3-2.7 (m, 6H), 3.2-3.4 (m, 1H), 3.7-4.0 (m, 4H), 8.18 (s, 2H); IR (KBr) ν: 3600-3100, 1610, 1555, 1510 cm⁻¹; UV (EtOH) λ: 244 (ε 19076), 330 (ε 1973);

Anal. Calcd. for C ₁₈ H ₂₉ N ₄ O·0.2H ₂ O:	
	C, 63.58; H, 8.71; N, 16.48.
Found:	C, 63.84; H, 8.71; N, 16.51.

EXAMPLE 23

1-(4-Fluoronaphth-1-yl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl]butanol

A solution of 1-(4-fluoronaphth-1-yl)-4-(5-fluoro-2-pyrimidinyl)piperazine (20 g, 5.0 mmol) in THF (20 mL) was treated with a solution of NaBH₄ (0.094 g, 2.5 mmol) in EtOH (20 mL) and stirred at 23 °C for 2 h. The reaction mixture was acidified to pH 1 with aqueous HCl solution (13 mL, 1N) then basified with aqueous NaOH solution (20%) to pH 9 and diluted with EtOAc (400 mL). The resulting mixture was washed with water (2 x 20 mL) and brine, dried (MgSO₄) and concentrated in *vacuo* to a crude material which was chromatographed on a silica gel pad (6.7 x 3 cm) using a mixture of 0-100% EtOAc in CH₂Cl₂. Evaporation of appropriate fractions gave 1.92 g (96%) of title compound. Recrystallization from EtOH gave an analytical sample, m.p. 129-31 °C; ¹H NMR (CDCl₃, 200 MHz) δ: 1.7-2.0 (m, 3H), 2.1-2.4 (m, 1H), 2.4-2.7 (m, 4H), 2.7-2.9 (m, 2H), 3.8-4.0 (m, 4H), 5.3-5.5 (m, 1H), 7.1-7.2 (m, 1H), 7.19 (s, OH, 1H), 7.4-7.8 (m, 3H), 8.0-8.3 (m, 4H), 8.2 (s); IR (KBr) ν: 3600-3310, 3300-3000, 1635, 1610, 1604, 1585, 1555, 1495 cm⁻¹; UV (CH₃CN) λ:

226 (€ 35991), 246 (€ 15791), 288 (€ 4801);

5

Anal. Calcd. for C ₂₂ H ₂₄ N ₄ OF ₂ :	
	C, 66.32; H, 6.07; N, 14.06.
Found:	C, 66.44; H, 6.09; N, 13.77.

10

EXAMPLE 24

15

 α -(4-Fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine pentanenitrile hydrochloride

20

25

To a mixture of potassium t-butoxide (5.28 g, 0.042 mole) in 1,2-dimethoxyethane (45 mL) cooled to -4° C was added dropwise over the course of 0.5 h a solution of 1-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazinebutan-1-one (6.21 g, 0.018 mole) and tosylmethylisocyanide (4.68 g, 0.024 mole) in 1,2-dimethoxyethane (75 mL) and EtOH (1.8 mL). After completion of addition, the mixture was stirred at 25° C for 0.5 h then heated at 45° C for 18 h. The reaction mixture was filtered, the filtrate concentrated in vacuo and the residue dissolved in EtOAc and flash-chromatographed on silica gel using EtOAc as eluant. Appropriate fractions were combined and concentrated in vacuo to afford 2.2 g (34%) of the product free base. The base was dissolved in EtOH (25 mL) and treated with ethanolic HCl to obtain the hydrochloride salt which was collected by filtration and dried affording 1.88 g, mp 234-236° C.

By following substantially the procedures described above in the description of the invention and in the above actual examples, additional Formula I compounds may be prepared. Some additional Formula I compounds are listed in Table 1.

30

EXAMPLE 25

35

Anoxic Nitrogen Test in Rats

40

The animals utilized are male Sprague-Dawley rats (200-240 grams) housed four animals per cage in a normal controlled environment with unlimited access to food and water. Usually there are 8 animals per dose, however, 4 animals can be employed to obtain an initial impression of a compound's activity. Animals surviving the anoxic insult are sacrificed via CO₂ inhalation following completion of the observation session (2 hr).

45

METHOD

50

Animals are parenterally or orally administered the vehicle or test compound 30 minutes prior to the anoxic insult. The anoxic episode consists of placing up to 4 animals in the sealed test chamber (10" 1 x 10" w x 6" h) continuously flushed with pure N₂ (4.5 grade) at a flow rate of 3 SCFM for 1 min. Animals are then promptly removed to normal atmosphere and observed for the 2 hour time period. Typically, animals become disoriented within 15 sec which leads into convulsions at 30 to 35 sec after which they remain motionless.

55

In spite of the fact that after the N₂ exposure the heart is still beating, all control animals fail to gasp when removed from the chamber and usually expire within 3 minutes. Drug treated animals, however, still gasp or start gasping after being removed which is a good indication if an animal will survive the N₂ exposure (1).

Results are recorded as:

Number of animals surviving (2 hr)Number of animals tested

and are statistically evaluated using the Finney Dose Response program for determination of the ED₅₀ and its corresponding 95% confidence limits.

DRUG EXAMPLE

10		# SURVIVING/#TESTED				
		Dose mg/kg, ip				ED ₅₀ (19/20 C.L)
		<hr/>				
15	Sabeluzole	<u>5.0</u> 3/8	<u>10</u> 15/16	<u>20</u> 18/20		4.7 (0.2-7.6)
	(+)MK-801	<u>0.5</u> 0/8	<u>1.0</u> 8/24	<u>2.0</u> 8/24	<u>4.0</u> 8/16 <u>10</u> 7/8	3.1 (2.0-6.9)

REFERENCES

1. Wauquier, A. et al; Arch. Int. Pharmacodyn., 249: 330-334 (1981).
2. Wauquier, A. et al; Drug Dev. Res., 8: 373-380 (1986).

Compounds of the present invention were rated at each dose level tested using the following rating scale:

- I = inactive (0% survival)
- + = weak activity (up to 25% survival)
- ++ = moderate activity (25-50% survival)
- +++ = good activity (51-75% survival)
- ++++ = very good activity (76-100%)


Table 2 contains test data for representative Formula I compounds. The highest rated dose level is the one displayed in Table 2.

Table 1

Additional Formula I Compounds

5

10



15

No.	Z	X	n	m	R ¹	R ²	R ³
-----	---	---	---	---	----------------	----------------	----------------

20

26.		CH ₂ NHCO-	3	0	H	F	H
-----	--	-----------------------	---	---	---	---	---

25

27.		CH ₂ NHCO ₂ -	3	0	H	Cl	H
-----	--	-------------------------------------	---	---	---	----	---

30

28.		CH ₂	3	0	H	F	H
-----	--	-----------------	---	---	---	---	---

35

40

29.		CH ₂ NHCO-	3	0	H	Br	H
-----	--	-----------------------	---	---	---	----	---

45

30.		CH ₂ NHCO ₂ -	3	0	H	F	H
-----	--	-------------------------------------	---	---	---	---	---

50

31.		CH ₂	3	0	H	F	H
-----	--	-----------------	---	---	---	---	---

55

Table 1 (cont'd)

Additional Formula I Compounds







No.	Z	X	n	m	R ¹	R ²	R ³
32.		CHOH	3	0	CH ₃	F	H
33.			3	0	H	F	H
34.		CH ₂	3	0	H	F	H
35.		CH ₂	3	0	H	F	H
36.		CHOH	2	1	H	F	H


Table 2

Ex. No.	Dose (mg/kg ip)	Rating
9	40	++
	40	+
	40	+++
	40	+
10	40 ^{a)}	+
11	40	+
21	40	+
18	40	+
19	40	++++
22	80	+++
20	80	++++
23	100	++++
13	40	++
12	80	++++
	40	+

a) given 60 minutes prior to anoxia testing.

Additional Detailed Description of The Invention


Some additional compounds related to Formula I have been prepared and found to have the useful anti-ischemic properties of the previous compounds of Formula I. These additional compounds extend the structural description of the X moiety in Formula I to give Formula I' which is the same as Formula I except for X.




Group X is now defined as a member selected from the group consisting of -O-, -S-, -SO₂-, -CO-;



wherein R⁴ is hydrogen or C₁₋₆ alkyl and R⁷ is hydrogen, C₁₋₆ alkyl, C₂₋₇ alkanoyl or



wherein W is hydrogen, halogen or alkoxy; and -CHR⁵- wherein R⁵ is hydrogen, CN, N₃ or NHR⁶ with R⁶ being R⁷ or



By C₁₋₆ alkyl (for R⁴), it is intended that both linear alkyl and cyclo alkyl moieties are included. By C₂₋₇ alkanoyl is intended alkylcarbonyl groups such as acetyl, propanoyl, cyclohexanoyl and the like.

These additional compounds were prepared by employing the general synthetic processes described hereinabove, using alterations which would be apparent to a skilled chemist in order to produce the desired product compound. Some additional examples are provided hereinbelow as guidance for synthesis of Formula I' compounds where X has been extended in structural definition

Compounds of Formula I'Example 371-Cyclohexyl-1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butanol

A cold (15° C) solution of 1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butanone (2.1 g, 6.1 mmol) in dry tetrahydrofuran (25 mL) kept under argon atmosphere was treated dropwise (15 min) with a solution of cyclohexylmagnesium chloride in ether (2.0 M, 3.18 mL, 6.36 mmol). The cooling bath was removed and reaction mixture after being stirred at 23° C for 2 h was treated dropwise (15 min) with HCl (2N, 3.5 mL), stirred for 15 min and diluted with CHCl₃ (50 mL). The organic phase was basified with

NaHCO₃ and separated. Aqueous phase was extracted with CHCl₃ (15 mL). The organic extracts were dried (MgSO₄) and concentrated in vacuo to a sticky solid which was purified on silica gel column using a mixture of 40-100% AcOEt in CHCl₃ as eluting solvent. The first group of fractions was concentrated in vacuo to give the carbinol as a solid, 1.1 g, 42%. The second group of fractions gave, after evaporation of solvent, the initial ketone 1.0 g, 48%. The carbinol was recrystallized from EtOH to give an analytical sample, mp 53-9° C.

Anal. Calcd. for C ₂₄ H ₃₂ F ₂ N ₄ O 0.25 C ₂ H ₆ O:	
Found:	C, 66.57; H, 7.64; N, 12.67. C, 66.37; H, 7.56; N, 12.53.

Anti-ischemic Rating + + + at 40 mg/kg ip.

Example 38

1-(4-Fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl acetate

A cold (5° C) mixture of 1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butanol monohydrochloride 1 (3.0 g, 7.8 mmol) and Et₃N (2.38 mL, 17.1 mmol) in CH₂Cl₂ (55 mL) was treated dropwise (10 min) with AcCl (0.67 mL, 9.4 mmol). The reaction mixture was stirred at 5° C for 0.25 h and then at 23° C for 2 h before being diluted with CH₂Cl₂ (250 mL). The organic solution was washed with water (30 mL), saturated Na₂CO₃ solution to bring pH to 9, water (20 mL) and brine, dried (MgSO₄) and concentrated in vacuo to a solid which was purified on silica gel column using a mixture of 30-40% AcOEt in CH₂Cl₂. The appropriate fractions were collected and concentrated in vacuo to a white solid, 2.93 g, 96%. Recrystallization from etherhexane mixture afforded analytical sample, mp 76-7° C.

Anal. Calcd. for C ₂₀ H ₂₄ F ₂ N ₄ O ₂ :	
Found:	C, 61.53; H, 6.20; N, 14.35. C, 61.55; H, 6.19; N, 14.33.

Example 39

1-(4-Fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl 4-fluorobenzoate hydrochloride

A cold (5° C) mixture of 1-(4-fluorophenyl)-4-(4-(5fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butanol monohydrochloride. (3.0 g, 7.8 mmol) and Et₃N (2.40 mL, 17.2 mmol) in CH₂Cl₂ (55 mL) was treated dropwise (10 min) with 4-fluorobenzoyl chloride (1.11 mL, 9.4 mmol). The cooling bath was removed and the reaction mixture stirred at 23° C for 20 h before being diluted with CH₂Cl₂ (250 mL). The organic solution was washed with water (30 mL), saturated Na₂CO₃ solution to bring pH to 9, water (30 mL) and brine, dried (MgSO₄) and concentrated in vacuo to a crude mixture which was purified on silica gel column using a mixture of 20-30% AcOEt in CH₂Cl₂. The appropriate fractions were concentrated in vacuo to a thick syrup, 3.4 g, 93%. The free base was solubilized in EtOH and treated with HCl in EtOH (one equivalent). The solvent was removed under reduced pressure leaving a gum which was crystallized from AcOEt affording

analytical sample mp 94-8° C.

Anal . Calcd. for C ₂₅ H ₂₅ F ₃ N ₄ O ₂ 1.15 HCl:	
	C, 58.60; H, 5.14; N, 10.93; Cl, 7.96.
Found:	C, 58.87; H, 5.17; N, 10.89; Cl, 7.94.

Anti-ischemic rating + + + at 40 mg/kg ip.

Example 40

1-(4-Azido-4-(4-fluoro-1-naphthyl)-1-butyl)-4-(5-fluoro-2-pyrimidinyl)piperazine monohydrochloride

To a cold (5° C) solution of Ph₃P (6.93 g, 26.4 mmol) in dry tetrahydrofuran (70 mL) kept under argon atmosphere was added dropwise (1 h) diisopropyl azodicarboxylate (5.2 mL, 26.4 mmol) and a solution of 1-(4-fluoro-1-naphthyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butanol 1 (9.2 g, 23.1 mmol) in dry tetrahydrofuran (140 mL), followed by the addition of a solution of diphenylphosphoryl azide (7.27 g, 26.4 mmol) in dry tetrahydrofuran (70 mL) over 10 min period. The reaction mixture was stirred at 5° C for 1 h then at 23° C for 2 h before being filtered. The cake was triturated several times in CH₂Cl₂ and filtered. All the filtrates were combined and concentrated in vacuo to a thick yellowish gum which was purified on silica gel column using a mixture of 0-30% AcOEt in CH₂Cl₂. The appropriate fractions were concentrated in vacuo leaving a colorless gum, 3.4 g, 35%. A solution of the free base in EtOH was treated with one equiv. of HCl in EtOH. The solution was concentrated in vacuo; the solid was crystallized from EtOH to give analytical sample, mp 212-3° C dec.

Anal . Calcd. for C ₂₂ H ₂₃ F ₂ N ₇ HCl:	
	C, 57.45; H, 5.26; N, 21.32.
Found:	C, 57.25; H, 5.05; N, 21.10.

Anti-ischemic rating + + at 40 mg/kg ip.

Example 41

1-(4-Fluoro-1-naphthyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl) butylamine hydrochloride

A mixture of 1-(4-azido-4-(4-fluoro-1-naphthyl)-1-butyl)-4-(5-fluoro-2-pyrimidinyl)piperazine 1 (0.79 g, 1.87 mmol) and 10% Pd/C (0.08 g) in EtOH (17 mL) was hydrogenated at 23° C for 3 h under 40 psi and then filtered. The filtrate was concentrated in vacuo to a gum which was purified on silica gel column using a mixture of 0-10% MeOH in CH₂Cl₂. The appropriate fractions were concentrated in vacuo giving a white solid; 0.50 g, 68%. A solution of the free base in EtOH was treated with HCl in EtOH (1 equiv); on standing, the HCl salt crystallized out; mp 228-30° C dec.

1. See Example 23 for synthesis.

1. See Example 40.

Anal . Calcd. for $C_{22}H_{27}F_2N_5$ 1.2 HCl:	
Found:	C, 59.89; H, 5.98; N, 14.87; Cl, 9.64. C, 59.95; H, 5.94; N, 15.75; Cl, 9.66.

5

Anti-ischemic rating + + + + at 40 mg/kg ip.

10

Example 42

1-[4-(4-Fluorophenyl)-4-methoxybutyl]-4-(5-fluoropyrimidin -2-yl) piperazine hydrochloride

15

A mixture of 1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butanol (4.79 g, 13.7 mmol) and sodium hydride (0.34 g, 14.2 mmol) in dry tetrahydrofuran was refluxed for 4 h under argon atmosphere before being cooled at 10 °C and treated with MeI (0.85 mL, 1.37 mmol). The reaction mixture was stirred at 25 °C for 24 h and diluted with CH_2Cl_2 (200 mL) and water (40 mL). The organic phase was separated and aqueous phase extracted with CH_2Cl_2 (10 mL). The organic extracts were dried ($MgSO_4$) and concentrated in vacuo to a crude mixture which was triturated in hexane and filtered. The filtrate was concentrated in vacuo and purified on silica gel column using AcOEt as eluting solvent. The appropriate fractions were concentrated in vacuo to a syrup, 3.3 g, 66%. The free base was solubilized in EtOH and resulting solution was treated with HCl (1.2 M, 1.2 equiv.) in EtOH. The solution was concentrated to dryness and the solid was recrystallized from EtOH; mp 199-201 °C.

25

Anal . Calcd. for $C_{19}H_{24}N_4OF_2$ 1.0 HCl:	
Found:	C, 57.21; H, 6.32; N, 14.05. C, 57.22; H, 6.32; N, 13.97.

30

35 Anti-ischemic rating + + at 40 mg/kg ip.

Additional Formula I' compounds are displayed in Table 3.


40

45

50

55




Table 3
Additional Formula I' Compounds



No.	Z	X	n	Yield%	MP(°C)	Anti-ischemic
						Rating
26			3	79	133-135	++
28		CH ₂	3	86	75-76	+++
30			3	63	123-125	++
31		CH ₂	3	32	72-73	++
43		CH2CH2COPh	3	77	165-166	++
44			3	82	178-179	++
45			3	76	130-131	+
46		CH2CH2CPh	3	92	159-160*	++
47		CH ₂	1	78	85-86	+
48		C=O	3	43	68-69	++++
49			3	17	142-144	+


1) Test compound given at a dose of 40 mg/kg i.p. 60 minutes prior to anoxia testing.

Table 3 (cont'd)
Additional Formula I' Compounds


5	<u>No.</u>	<u>Z</u>	<u>X</u>	<u>n</u>	<u>Yield%</u>	<u>MP (°C)</u>	Anti-ischemic <u>Rating</u>
10	50			3	93	165-166	-
	51		CHNH ₂	3	52	277-8B*	-
15	*Mp of hydrochloride salt						

Claims

1. A compound of Formula I' and its pharmaceutically




acceptable acid addition salts and/or solvates thereof wherein
Z is a member selected from the group consisting of




naphthalenyl, anthracenyl, fluorenyl, phenanthrenyl, and C₅₋₆ cycloalkyl;


X is a member selected from the group consisting of -O-, -S-, -SO₂-, -CO-



wherein R⁴ is hydrogen, or C₁₋₆ alkyl and R⁷ is hydrogen, C₁₋₆ alkyl, C₂₋₇ alkanoyl, or



wherein W is hydrogen, halogen or alkoxy, and -CHR⁵-wherein R⁵ is hydrogen, CN, N₃ or NHR⁶ with R⁶ being R⁷ or



5 or Z and X taken together can be



10 R¹ is hydrogen or C₁₋₄ alkyl;
 R² is halogen;
 R³ is hydrogen, C₁₋₄ alkoxy or C₁₋₄ alkylthio;
 n is 1 - 3; and
 15 m is 0 or 1; with the proviso that Z cannot be



20 when X is or



25

or -CO-, while R³ is either hydrogen or C₁₋₄ alkoxy, or while m is 0.

2. A compound of claim 1 wherein Z is



30

3. A compound of claim 1 wherein X is -CHR⁵-.

4. The compounds of claim 1:

35 4-[5-fluoro-4-(methylthio)-2-pyrimidinyl]-α-(4-fluorophenyl)-1-piperazinebutanol;
 α-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazinepentanenitrile;
 1-(4-fluorophenylbutyl)-4-(5-fluoro-2-pyrimidinyl)piperazine;
 1-[3-(4-fluorophenoxy)propyl]-4-(5-fluoro-2-pyrimidinyl)-piperazine;
 1-[3-(4-fluorophenylthio)propyl]-4-(5-fluoro-2-pyrimidinyl)-piperazine;
 40 1-[3-(1,3-dioxolan-2-yl) propyl]-4-(5-fluoro-2-pyrimidinyl)piperazine;
 1-[3-(4-fluorophenylsulfonyl) propyl]-4-(5-fluoro-2-pyrimidinyl)piperazine;
 1-[4-acetamido-4-(4-fluorophenyl)butyl]-4-(5-fluoro-2-pyrimidinyl) piperazine ;
 1-[4-(4-fluorobenzanido)-4-(4-fluorophenyl)butyl]-4-(5-fluoro-2-pyrimidinyl) piperazine;
 1-cyclohexyl-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl]butanol;
 45 1-[4-(4-fluorophenyl)-4-(4-fluorophenylsulfonamidol)butyl]-4-(5-fluoro-2-pyrimidin -yl)-piperazine ;
 1-(4-fluoronaphthyl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl]butanol ;
 1-(4-fluoronaphthyl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl]butanone ;
 4-[5-(5-fluoro-2-pyrimidinyl)(1R,4R)-2,5-diazabicyclo[2.2.1] heptan-2-yl]-1-(4-fluorophenyl)butanol;
 4-[5-(5-fluoro-2-pyrimidinyl)(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-1-(4-fluorophenyl)butanol;
 50 1-(2-thienyl)-4-[4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl]butanone ;
 N-(1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl) benzamide ;
 1-cyclohexyl-1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butanol;
 1-(4-cyclohexyl-1-butyl)-4-(5-fluoro-2-pyrimidinyl)piperazine;
 N-(1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl 4-fluoro-1-naphthamide;
 55 N-(1-(4-fluoro-1-naphthyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl) 4-fluorophenylsulfonamide ;
 1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl acetate;
 1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl benzoate ;
 1-(2-(4-fluorophenyl)-1-ethyl) -4-(5-fluoro-2-pyrimidinyl)piperazine;


- 1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl-4-fluorobenzoate ;
 1-(4-(4-fluoro-1-naphthyl)-1-butyl)-4-(5-fluoro-2-pyrimidinyl)piperazine ;
 1-(4-azido-4-(4-fluoro-1-naphthyl)-1-butyl)-4-(5-fluoro-2-pyrimidinyl)piperazine ;
 1-cyclohexyl-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butanone ;
 5 N-(1-(4-fluoro-1-naphthyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl)-4-fluorobenzamide ;
 N-(1-cyclohexyl-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl)-4-fluorophenyl-sulfonamide ;
 N-(1-cyclohexyl-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl)-4-fluorobenzamide ;
 1-(4-fluoro-1-naphthyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butylamine ;
 N-(1-(4-fluorophenyl)-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butyl)-cyclohexane-carboxamide ;
 10 1-cyclohexyl-4-(4-(5-fluoro-2-pyrimidinyl)-1-piperazinyl)-1-butylamine ;
 1-[4-(4-fluorophenyl)-4-methoxybutyl]-4-(5-fluoropyrimidin-2-yl) piperazine.

5. The use of at least one compound of anyone of claims 1 to 4 for preparing a pharmaceutical composition for therapeutically or prophylactically protecting brain cells from ischemia.

6. A pharmaceutical composition suitable for systemic administration to a mammalian host comprising a pharmaceutical carrier and at least one compound of anyone of claims 1 to 4.

7. A process for preparing the pharmaceutical composition of claim 6 which comprises incorporating at least one compound of anyone of claims 1 to 4 into a pharmaceutical carrier.

8. A process for preparing the compounds of claims 1 to 4 which comprises reacting a compound of the general formula II:




II

wherein R¹, R², R³ and m are as defined in claim 1, with a compound of the general formula III:

Z-A-(CH₂)_n-W

- 30 wherein Z and n are as defined in claim 1, A represents O, S, CO and CH₂ and W represents a leaving group, to obtain a compound of the general formula IA:



Ia

40 and, if desired, converting a compound of formula Ia to the other compounds of formula I by transformation of the A moiety.



EP 90110399.4

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.)
X	DE - A1 - 2 319 016 (BUZAS) * Formula II *	1,6	C 07 D 403/04 C 07 D 409/04 C 07 D 409/14 C 07 D 239/42
X	DE - A1 - 2 053 759 (SUMITOMO) * Formula I, II; table III; compound 6 *	1,6	A 61 K 31/505
X	DE - A1 - 3 507 983 (KRISTOL) * Compound 1c *	1,6	
D, X	US - A - 4 605 655 (YEVICH) * Compound 16 *	1,6	
D, X	US - A - 2 973 360 (JANSSEN) * Example 4 *	1,6	
A	WO - A1 - 88/07 528 (AKTIEBOLOGET LEO) * Claim 1 *	1,6	TECHNICAL FIELDS SEARCHED (Int. Cl.)
A	DE - A1 - 3 339 943 (SANDOZ) * Claims 1,9 *	1	C 07 D 403/00 C 07 D 409/00 C 07 D 239/00
A	EP - A1 - 0 183 191 (YOSHITOMI) * Claim 1 *	1	
A	CHEMICAL ABSTRACTS, vol. 109, no. 7, August 15, 1988, Columbus, Ohio, USA KAWAKITA, Takeshi et al. "Preparation of ((heterocycly-alkyl)thienyl)thiazole derivatives as drugs"	1	
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 27-08-1990	Examiner HAMMER
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.)
A	<p>page 687, column 1, abstract- -no: 54 766s & Jpn. Kokai Tokkyo Koho JP 62,192,379 (87,192,379) --</p> <p>CHEMICAL ABSTRACTS, vol. 94, no. 1, January 5, 1981, Colum- bus, Ohio, USA PESSON, MARCEL et al. "Synthetic antibacterial agents. Pipemidic acid derivatives" page 372, column 2, abstract- -no. 3 981x & Eur. J. Med. Chem. - Chim. Ther. 1980, 15(3),263-8 (Fr.) --</p>	1	
A	<p>CHEMICAL ABSTRACTS, vol. 110, no. 7, February 13, 1989, Columbus, Ohio, USA MC MILLER, BRIAN A. et al. "Reversal of neuroleptic- -induced catalepsy by novel arylpiperazine anxiolytic drugs" page 63, column 1, abstract- -no. 51 191k & J. Pharm. Pharmacol. 1988, 40(12), 885-7 (Eng) ----</p>	1,6	
			TECHNICAL FIELDS SEARCHED (Int. Cl.)
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 27-08-1990	Examiner HAMMER
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.